

Gravitational Collapse End States

Pankaj S Joshi

Department of Astronomy and Astrophysics

Tata Institute of Fundamental Research

Homi Bhabha Road, Mumbai 400 005, India

Recent developments on the final state of a gravitationally collapsing massive matter cloud are summarized and reviewed here. After a brief background on the problem, we point out how the black hole and naked singularity end states arise naturally in spherical collapse. We see that it is the geometry of trapped surfaces that governs this phenomena.

It was pointed out by S. Chandrasekhar in 1935:

”...The life history of a star of small mass must be essentially different from that of a star of large mass... A small mass star passes into White-dwarf stage... A star of large mass *cannot* pass into this stage and one is left speculating on other possibilities.”

The question that *what happens when a star dies* has been a key problem in astronomy and astrophysics for past decades. If the star is sufficiently massive, beyond the white dwarf or neutron star mass limits, then a continued gravitational collapse must ensue when the star has exhausted its nuclear fuel.

What are the *possible end states* of such a *continued gravitational collapse*? To answer this question, one must study *dynamical collapse scenarios* within the framework of a gravitation theory such as Einstein’s theory. I shall try to give a somewhat non-technical account of recent developments in this area in recent years.

Penrose conjectured in 1969, that the ultra-dense regions i.e. the spacetime singularities (where the physical quantities e.g. densities, curvatures are having extreme values) forming in gravitational collapse must be hidden within the event horizon of gravity, that is, the collapse must end in a black hole. This is called the Cosmic Censorship Conjecture. There is however no proof, or any suitable mathematical formulation of the same, available as of today.

Under the situation, very many researchers have made extensive study of various dynamical collapse models, mainly spherically symmetric, over past decade and a half, to investigate the final outcome of an endless gravitational collapse. When no proof, or even a suitable mathematical formulation of censorship conjecture is available, it is only such studies that can throw light on this

issue. The generic conclusion appears to be: Either a Black Hole(BH) or a Naked Singularity(NS) develops as end product of collapse, depending on the initial data (e.g. initial density, pressures, and velocity profiles for the collapsing shells) from which the collapse develops, and the nature of dynamical evolutions as permitted by Einstein equations.

While extensive study is made of astrophysics of Black Holes, for NS we may still want to inquire into questions such as, *Are Naked Singularities Generic*, or *What are the Physical factors* that cause NS, rather than a BH forming as Collapse End State? That is, one may wish to understand in a better way the naked singularity formation in gravitational collapse. We try to address these issues in some detail here, and this may also possibly give an idea on the present status of Cosmic Censorship Conjecture. For some further details and references, we refer to [1-6].

A spherically symmetric spacetime with the general metric in t, r, θ, ϕ co-ordinates can be written as,

$$ds^2 = -e^{2\nu(r,t)}dt^2 + e^{2\psi(r,t)}dr^2 + R^2(t,r)d\Omega^2 \quad (1)$$

In such a comoving reference frame, the energy-momentum tensor for a general ‘*Type I*’ matter field can be written as,

$$T_t^t = -\rho; \quad T_r^r = p_r; \quad T_\theta^\theta = T_\phi^\phi = p_\theta \quad (2)$$

This includes practically all physically reasonable matter forms such as dust, perfect fluids, massless scalar fields, and such other forms of matter.

We always consider the matter to satisfy a suitable energy condition, Such as the *Weak Energy Condition*:

$$T_{ik}V^iV^k \geq 0 \quad (3)$$

The point here is that, while it may be possible that energy condition violations may occur in the very late stages of collapse when the quantum effects take over, they should necessarily be satisfied for the classical phase which governs most of the collapse evolution.

The *Einstein equations* then have the following form:

$$\rho = \frac{F'}{R^2 R'}; \quad p_r = \frac{-\dot{F}}{R^2 \dot{R}} \quad (4)$$

$$\nu' = \frac{2(p_\theta - p_r)}{\rho + p_r} \frac{R'}{R} - \frac{p_r'}{\rho + p_r} \quad (5)$$

$$-2\dot{R}' + R'\frac{\dot{G}}{G} + \dot{R}\frac{H'}{H} = 0 \quad (6)$$

$$G - H = 1 - \frac{F}{R}$$

The functions G and H above are defined as,

$$G(t, r) = e^{-2\psi}(\dot{R}')^2, H(t, r) = e^{-2\nu}(\dot{R})^2 \quad (7)$$

Also $F = F(t, r)$ is an arbitrary function, and has the interpretation of the mass function of the collapsing cloud, with $F \geq 0$. Various *Regularity Conditions* ensure that the collapse develops from a Regular Initial Data. From the point of view of dynamic evolution of initial data, at the initial epoch $t = t_i$, we have six arbitrary functions of the radial coordinate r ,

$$\begin{aligned} \nu(t_i, r) &= \nu_0(r); \quad \psi(t_i, r) = \psi_0(r); \quad R(t_i, r) = R_0(r) \\ \rho(t_i, r) &= \rho_0(r); \quad p_r(t_i, r) = p_{r_0}(r); \quad p_\theta(t_i, r) = p_{\theta_0}(r) \end{aligned} \quad (8)$$

All the initial data represented above are not mutually independent, as from *Bianchi Identity* we have,

$$\nu_0(r) = \int_0^r \left(\frac{2(p_{\theta_0} - p_{r_0})}{r(\rho_0 + p_{r_0})} - \frac{p'_{r_0}}{\rho_0 + p_{r_0}} \right) dr$$

Hence we see that there are five total field equations with seven unknowns, ρ , p_r , p_θ , ψ , ν , R , and F , giving us the freedom of choice of two free functions. Selection of these functions, subject to the weak energy condition, and the given initial data for collapse at the initial spacelike surface, determines the matter distribution and metric of the space-time, and thus leads to a particular time evolution of collapse from the given initial data. Let us now consider a general mass function $F(t, r)$ for the collapsing cloud, given as

$$F(t, r) = r^3 \mathcal{M}(r, v) \quad (9)$$

where $\mathcal{M}$ is any regular and suitably differentiable function. We note here that we are NOT making any choice regarding the mass function, as it is in the most general form.

Using the freedom of choice mentioned above, let us choose,

$$\nu(t, r) = A(t, R) \quad (10)$$

(assuming that there are no shell-crossings in the spacetime, one could also make a more general choice as defined by $\nu(t, r) = A(r, v)_v R$). Then we get the solution to *Einstein equation* as,

$$G(t, r) = b(r)e^{2(A - \int A_t dt)} \quad (11)$$

Here $b(r)$ is another arbitrary function of r . In correspondence with dust models, we write,

$$b(r) = 1 + r^2 b_0(r) \quad (12)$$

where $b_0(r)$ is the velocity distribution function of the collapsing shells. The marginally bound case corresponds to $b_0(r) = 0$.

It is now possible to go back to the Einstein equations and work out the function $t(v, r)$. The physical spacetime singularity develops at the value $R = 0$, which corresponds to the physical radius of all collapsing shells going to zero. This corresponds to the time $t_s(r) = t(0, r)$. At the initial epoch we have $v = 1$ and the singularity is defined by the value $v = 0$. Expanding the function $t(v, r)$ around the center, we get,

$$t(v, r) = t(v, 0) + r\mathcal{X}(v) + O(r^2) \quad (13)$$

We then get the tangent to the singularity curve at the central singularity as,

$$\mathcal{X}(0) = -\frac{1}{2} \int_0^1 dv \frac{\sqrt{v}(b_1 v + v h_1(v) + \mathcal{M}_1(v))}{(b_{00} v + v h_0(v) + \mathcal{M}_0(v))^{\frac{3}{2}}} \quad (14)$$

To decide the final fate of collapse in terms of either a Black Hole or a Naked Singularity, we need to study the behavior of the apparent horizon in the spacetime, and to examine if there are any families of outgoing non-spacelike trajectories, which terminate in the past at the singularity.

The *apparent horizon* within the collapsing cloud is given by $R/F = 1$, which gives the boundary of the trapped surface region of the space-time. If the neighborhood of the center gets trapped earlier than the singularity, then it will be covered. Otherwise, the singularity could be naked with non-spacelike future directed trajectories escaping from it to outside observers.

In order to consider the families of outgoing null geodesics from the singularity, and to examine the nature of the central singularity at $R = 0$, $r = 0$, let us consider the outgoing radial null geodesics equation,

$$\frac{dt}{dr} = e^{\psi - \nu} \quad (15)$$

The singularity occurs at $v(t_s(r), r) = 0$. Therefore, if there are any future directed null geodesics terminating in the past at the singularity, we must have $R \rightarrow 0$ as $t \rightarrow t_s$ along these curves.

Rewriting the null geodesic equation in terms of the variables $(u = r^\alpha, R)$, it can be further analyzed,

$$\frac{dR}{du} = \frac{1}{\alpha} r^{-(\alpha-1)} R' \left[1 + \frac{\dot{R}}{R'} e^{\psi-\nu} \right] \quad (16)$$

Choosing $\alpha = \frac{5}{3}$ we get

$$\frac{dR}{du} = \frac{3}{5} \left(\frac{R}{u} + \frac{\sqrt{v}v'}{\sqrt{\frac{R}{u}}} \right) \left(\frac{1 - \frac{F}{R}}{\sqrt{G}[\sqrt{G} + \sqrt{H}]} \right) \quad (17)$$

The tangent to the null geodesics from the singularity is,

$$x_0 = \lim_{t \rightarrow t_s} \lim_{r \rightarrow 0} \frac{R}{u} = \left. \frac{dR}{du} \right|_{t \rightarrow t_s; r \rightarrow 0} \quad (18)$$

Using the null geodesic equation the value x_0 is given as,

$$x_0^{\frac{3}{2}} = \frac{5}{\sqrt{6}} \mathcal{X}(0) \quad (19)$$

Then, whenever $\mathcal{X}(0) > 0$, there is radial null geodesic coming out from the singularity as given by $t - t_s(0) = x_0 r^{\frac{5}{3}}$.

It has been already stated that we have the choice of two free functions in the above analysis, among which we are choosing only one so far, in the most general case, namely, the form of metric function $\nu(t, r)$. Therefore, the analysis holds for any equation of state of the form

$$p_r = Q(\rho) \quad (20)$$

In this case there would be a constraint on otherwise arbitrary function $\mathcal{M}$. Furthermore the value of the function $p_\theta(t, r)$ in terms of $\rho(t, r)$ will be determined from Einstein equations.

In other words, it turns out that the BH/NS phases are determined by the geometry of the trapped surfaces and the apparent horizon that develop as the collapse evolves.

What governs the geometry of the trapped surfaces, or the formation or otherwise of the naked singularities in the spacetime? In other words, what is it that causes the naked singularity to develop rather than a black hole as the final end product of collapse? It turns out that physical agencies such as INHOMOGENEITIES in matter profiles as well as spacetime SHEAR play an important role to distort the trapped surface geometry, and could delay the trapped surface formation during the collapse, thus giving rise to NS, rather than a BH, as collapse end state (see e.g. references [7-11]).

This, in a way, provides the physical understanding of the phenomena of BH/NS end states.

It is seen that in spherical collapse the BH/NS phases as end state of collapse are generic, and are seen to be determined by the nature of the initial data from which the collapse develops, and in terms of the allowed dynamical evolutions. Physical agencies such as Inhomogeneities and Shear cause them, and given the initial data, there are non-zero measure classes of evolutions that evolve into either of these outcomes.

What does the Non-spherical Collapse do? There are some examples which indicate outcome to be similar—but the evidence in that case is somewhat limited so far. The main difficulty is the complexity of the Einstein system, which is second order, non-linear partial differential equations.

There are several quite interesting questions which are under active investigation at the moment. For example, could naked singularities generate bursts of Gravity Waves? What kind of quantum effects will take place near NS? What will be the generic outcome for the case of non-spherical collapse? Many of these issues would have interesting physical implications. It appears quite likely from the current investigations that the astrophysical phenomena such as the Gamma Rays Bursts will have a strong connection to the physics and dynamics of gravitational collapse of massive stars. Another intriguing possibility could be, NS may possibly provide us with some kind of observational signatures for the Quantum Gravity effects from ultra-strong gravity regions. This last possibility would then be considered quite an exciting prospect.

-
- 1 A. Krolak, Prog. Theor. Phys. Suppl. **136**, 45 (1999).
 - 2 R. Penrose, in *Black holes and relativistic stars*, ed. R. M. Wald (University of Chicago Press, 1998).
 - 3 P. S. Joshi, Pramana **55**, 529 (2000).
 - 4 M. Celerier and P. Szekeres, Phys.Rev. D65 (2002) 123516.
 - 5 R. Giambo', F. Giannoni, G. Magli, P. Piccione, Commun.Math.Phys. 235 (2003) 545-563.
 - 6 T. Harada, H. Iguchi, and K. Nakao, Prog.Theor.Phys. 107 (2002) 449-524.
 - 7 P. S. Joshi, N. Dadhich and R. Maartens, Phys. Rev **D65**, 101501R (2002).
 - 8 E. Brinis, S. Jhingan and G. Magli, Class. Quant. Grav. **17**, 4481 (2000).
 - 9 S. M. C. V. Goncalves, gr-qc/0312123.
 - 10 F. C. Mena, B. C. Nolan, and R. Tavakol, gr-qc/0405041.
 - 11 P. S. Joshi, R. Goswami and N. Dadhich, Phys. Rev. *D70*, 087502 (2004).